

# STUDY ON PRACTICABILITY OF HYDROPONICAL CULTURE OF GRAM (*CICER ARIETINUM*)

L. Prabhas<sup>1</sup>, Agrawal M<sup>2</sup>, Tamrakar V<sup>3</sup>

## ABSTRACT

Agriculture is a chief supplier for the growth of Indian Economy. Rice, Wheat, Peas, pulses and vegetables are some major plant species cultivated primarily. Due to rise in population and contradictory decline in land size available for agriculture, it is necessary to introduce a new method of agriculture for increasing potential of cultivation. Gram (*Cicer arietinum*) is one of them, also commonly known as chick pea. Hydroponical culture of chick pea plant showed some possibility for large scale production. Seed germination taken very less time in comparison to soil germinated seed. Growth of shoot was also measured after every 24hr, which illustrated elevated shoot length (cm) in short time period. Leaf initiation was also observed after 48hr of seed propagation. The feasibility of the technique for the selected plant species was predictable in liquid nutrient medium. Fruiting and flowering was not observed in the experiment and required some more attempts with modification in experimental design. Temperature, Humidity, light and other physical factors were maintained with natural surroundings.

**Keywords:** Hydroponics, Seed germination, Shoot length, Leaf initiation, propagation, Liquid Nutrient Medium.

## I. INTRODUCTION

A big part of world population relies on agriculture development for their economy and growth. We all are well aware of importance of agricultural products in human life. Advancement of human life and shortage of agricultural land is a serious threat for future food security. Advancement in science and technology is playing a major role of therapeutic for healing problem of food security since last few decades. Somehow literature revealed that in some part of the world a technique named as "Hydroponics" is contributing a significant role in enhanced agriculture productivity (Craver and Williams, 2014; Das and Sing Majhi, 2009; Flores-López et al. 2009; Hoagland and Arnon, 1950; Koohakan, 2008). According to the reviewed literature this technique is significantly useful to avoid effect of many environmental factors at large scale. This is the technique to grow plant in controlled environment. Environmental factors like temperature, humidity, light intensity etc. can be maintained or manipulated according to the need of selected plant species. Hence decrease in final productivity due to adverse effect of environmental factor can be avoided (Ruiz and Taleisnik, 2013; Smeets et al., 2008).

In this study small experimental setup is used to determine the practicability of Hydroponical technique for agricultural crop. Chick pea (*Cicer arietinum*) commonly known as Gram or Chana, is selected as modal plant. This plant belongs to family Fabaceae, subfamily Faboideae categorized in shrub. Generally average height of

plant is about 20-55 cm. Two common types of chick peas are grown and available in all part of the India. These are Desi Chana and Bombay chickpeas. Early literature revealed physical characteristics of the selected plant species seems fruitful the proposed experiment in initial circumstances. Indian chickpea is very popular among the country people because of its nutrient richness. Chick pea is a rich source of protein, carbohydrates, lipid content, fibre and minerals i.e. iron, magnesium, phosphorus, zinc etc. Amino acids such as tryptophan, lysine, isoleucine etc. are abundant in the selected plant product. With high calorific value selected plant species is proved to be unsurpassed for the experiment.

## II. MATERIAL AND METHOD

**Establishment of Hydroponic System:** Design of the Hydroponic System used for the experimentation was influenced by various source of research work done earlier through various novelist. (Du Toit, 2007; Ortiz et al., 2009; Kratky and Bowen, 1988; Roberto, 2000).

A miniature setup built for the accomplishment of experimental protocol. All the necessary articles were provided simultaneously. Some important features of the experimental setup has been discussed below;


**Nutrient Reservoir:** It is a collection and feeding unit for the model plant. All the necessary nutrients were dissolved in double distilled water in specific amount. Nutritional requirement of various organic and inorganic matters, such as Nitrogen (N), Carbon (C), Sulphur (S), Phosphorus (P), Potassium (K) etc. may differ in various plant species. This study may also demonstrate about stipulation of proper nutritional supply in plant species and the effect of concentration of various nutritional sources in plant species (Gericke, 1937; Shinohara et al., 2011; Yan et al., 2012; Asao, 2013). Non-reactive material plastic material is used to construct Nutrient reservoir. Continuous flow of nutrients dissolved in water is provided but rate of flow is controlled by valve system. Sequentially concentration of dissolved nutrient is checked after precise time interval.

**Temperature and Humidity:** With the help of thermostat, standard value of temperature for plant growth between 30-35°C was maintained through air-conditioning unit. Room temperature outside environment was approximately 40-45°C during the study period in the month of June. Humidity was maintained up to 95% which is optimum for plant growth. Apical necrosis, blackening and other vital diseases are very common in plants due to temperature fluctuation above and below specific optimum value suitable for plant species.

**Photoperiod and Light Intensity:** It is well known subject that plant growth is affected by the light spectra. Optimum light intensity was achieved from maintaining common range of color spectra. Earlier, it has been demonstrated that red (600-800 nm) and blue light (380-480 nm) are most favorable for plant growth (Caruso et al., 2011). In the present study, light spectra from 400 to 700 nm were supplemented through LED on hydroponically cultivated selected shrub. **Photosynthetically active radiation**, often abbreviated as **PAR**, designates the spectral range (wave band) of solar radiation from 400 to 700 nm that photosynthetic organisms use to optimize the process of photosynthesis. During the experimentation neutral photoperiod of about (LD 12:12) was preserved (Carre, 2001; Martineau; 2012; Thomas, 2006).

**Aeration:** Aeration in nutrient solution was achieved by the use of aerator (pump). This is commonly available in market, specially used in aquariums. The aeration pump helps to grab atmospheric air and mix it to nutrient

solution. Maintenance of proper aeration helps to improve plant growth (Von-Bieberstein, 2014; Souret and Weather, 2000; Park and Kurata, 2009).



### III. RESULT AND DISCUSSION

Hydroponical culture of chick pea plant showed some possibility for large scale production. In the beginning 50 chickpea seeds were propagated. Seed germination taken very less time in comparison to soil germinated seed. First significant physiological change in sowed seed was observed within 7hr. Growth of shoot was also measured after every 24hr, which illustrated elevated shoot length (cm) in short time period. It was about 1cm long structure with two cotyledons. Leaf initiation was also observed after 48hr of seed propagation (Wang, 1999). Root growth was also noteworthy, approximately 80% seeds showed root growth after 24hrs (Trang, 2010; Girdthai, 2010). The feasibility of the technique for the selected plant species was predictable in liquid nutrient medium. Fruiting and flowering was not observed in the experiment and required some more attempts with modification in experimental design. In aspect of total mass productivity it was found that the feasibility of the technique with selected plant species was good.

Some other characteristic for the accomplishment of the technique are following:

1. It was helpful to reduce reliance on climatic factors such as light, temperature and humidity.
2. Low land area with high mass productivity.
3. Reduction in soil born microbial and viral diseases.
4. Reduction in insect growth, helps to prevent use of harmful insecticides
5. Indoor hydroponical system helps to avoid productivity loss due to grazing.
6. Less water requirement: solution for irrigation.
7. There is no space for weed. Hence, prevention of use of weedicides.

### 3.1 Limitation

1. This new emerging technology is not suitable for the crops like rice, wheat etc. in which large field area is required.
2. This technique is most suitable for small herb, shrubs and climbers. Cultivation of trees and tall plants are not easy in hydroponics.
3. Proper knowledge and continuous safeguarding is required.
4. First time investment cost is higher than the cost of traditional technique.
5. Setup requires some scientific equipment and handling knowledge.

### 3.2 Future aspects

Future aspects from the mentioned technique are really very high. Pharmaceutical industries are facing problem of continuous supply of medicinal plant and their products. Many herbs and shrubs are known for their medicinal properties can be easily grown by hydroponic without altering their biochemical composition. Green house facility prevents the supply of plant and their product in specific season. In controlled environment we can cultivate plant throughout the year.

According to literature reviewed one of the best future aspects associated with hydroponics is space technology. It is well known practice that the space mission includes a long term accommodation at space station. It is really a big trouble to carry plenty of food products with many numbers of people in single spacecraft. Spoilage and maintenance of food product is also a significant problem in space missions. Hence, the scientists are trying to find out some substitute. Theoretically in place of large mass of food product, it is easy to carry seed and nutrients dissolved in water. Cultivation of plant, utilization and release of waste material in space eradicate burden from the spacecraft. We can reduce burden of weight in spacecraft in both the ways i.e. departure and arrival.

Indoor home gardening may be encouraged by this unique technique of horticulture. Many ornamental plant species are used to grow in indoor atmosphere frequently by the people. But the knowledge of plant cultivation with nutrient water medium will encourage them to cultivate economically important plant species. Not at very large scale but somehow slight load from the shoulder of farmer would be condensed inside home environment of the society. Desert and land with low water availability can also be used for horticulture activity with minimum requirement of water and nutrient sources (Thompson, 1978) .

Some of the earlier studies also showed vivid future possibilities in molecular biology. Study based on molecular marker through hydroponics and effect of various chemical and environmental factors on Genetic composition in hydroponical culture has been well illustrated (Tuberosa et al., 2002). Bioremediation through hydroponics is also possible via cultivation of various plant species. Many plant species are able to utilize and absorb harmful heavy metals and chemicals from water for their growth and development. Those heavy metals and chemicals are dreadful for human health (Singh et al., 2012; Singh, 2010; Snow, and Ghaly, 2008; Rababah and Al-Shuha, 2009; Paz-Alberto and Sigua, 2013).

## IV. ACKNOWLEDGEMENT

We would like to express my sincere thanks to all the faculty members and staff member of the school who provide us all the necessary support to do this wonderful research activity. We came to know about lots of new things we are really thankful to them. Secondly we are also grateful to all the people directly or indirectly involved in this action of learning. We hope that we shall have such support, encouragement and inspiration in our future activity too.

## Bibliography

- [1] Albery, W. J., Haggett, B. G. D., and Svanberg, L. R. (1985). The development of sensors for hydroponics. *Biosensors*, 1(4), 369-397.
- [2] Asao, T., Shimizu, N., Ohta, K., and Hosoki, T. (1999). Effect of rootstocks on the extension of harvest period of cucumber (*Cucumis sativus* L.) grown in non-renewal hydroponics. *Journal of the Japanese Society for Horticultural Science*, 68(3), 598-602.
- [3] Akyas, A. M., Widayat, D., and Nursuhud. (2005). Research and development in hydroponics technology at the laboratory of horticulture padjadjaran university (A case with tomato cuvar. recento). *Journal of Agriculture and Rural Development in the Tropics and Subtropics Supplement*, (83), 87-96.
- [4] Ako, H., and Baker, A. (2009). Small-Scale Lettuce Production with Hydroponics or Aquaponics. *Sustainable Agriculture*, 2, 1-7
- [5] Al-Khalifah, N. S., Shanavaskhan, A. E., and Khan, F. A. (2010). Utilizing hydroponics technique for acclimatizing tissue culture derived plantlets under desert environment. *Acta Horticulturae*, 865, 163-170
- [6] Asao, T., Asaduzzaman, M., Mondal, M. F., Tokura, M., Adachi, F., Ueno, M., and Ban, T. (2013). Impact of reduced potassium nitrate concentrations in nutrient solution on the growth, yield and fruit quality of melon in hydroponics. *Scientia Horticulturae*, 164, 221-231.
- [7] Brown, C. S., Cox, W. M., Dreschel, T. W., and Chetirkin, P. V. (1992). The vacuum-operated nutrient delivery system: Hydroponics for microgravity. *HortScience*, 27(11), 1183-1185.
- [8] Bianchi-Hall, C. M., Carter Jr., T. E., Bailey, M. A., Mian, M. A. R., Rufty, T. W., Ashley, D. A., and Parrott, W. A. (2000). Aluminum tolerance associated with quantitative trait loci derived from soybean PI 416937 in hydroponics. *Crop Science*, 40(2), 538-545.
- [9] Carre I. A. (2001). Day-Length perception and the photoperiodic regulation of flowering in *Arabidopsis*. *Journal of Biological Rhythms*, 16(4), 415-423.
- [10] Caruso, G., Villari, G., Melchionna, G., and Conti, S. (2011). Effects of cultural cycles and nutrient solutions on plant growth, yield and fruit quality of alpine strawberry (*Fragaria vesca* L.) grown in hydroponics. *Scientia Horticulturae*, 129(3), 479-485.
- [11] Craver, J. K., and Williams, K. A. (2014). Assessing student learning from an experiential module in a greenhouse management course using hydroponics and recirculating solution culture. *HortTechnology*, 24(5), 610-617.

- [12] Da Silva, A. O., Soares, T. M., e Silva, Ê. F. F., dos Santos, A. N., and Klar, A. E. (2012). Rucola water consumption in NFT hydroponics using wastewater desalinator in ibimirim, PE (brazil). *IRRIGA*, 17(1), 114-125.
- [13] Du Toit, A. G. A., and Labuschagne, M. T. (2007). A comparison between hydroponics systems and pots for growing wheat in the greenhouse. *South African Journal of Plant and Soil*, 24(2), 120-123.
- [14] Das, A., and Sing Majhi, D. (2009). Low cost hydroponics devices and use of harvested water for vegetable and flower cultivation. *Journal of Applied Horticulture*, 11(2), 123-126.
- [15] Dos Santos, J. D., Lopes da Silva, A. L., da Luz Costa, J., Scheidt, G. N., Novak, A. C., Sydney, E. B., and Soccol, C. R. (2013). Development of a vinasse nutritive solution for hydroponics. *Journal of Environmental Management*, 114, 8-12.
- [16] Flores-López, R., Sánchez-Del Castillo, F., Rodríguez-Pérez, J. E., Colinas-León, M. T., Mora-Aguilar, R., and Lozoya-Saldaña, H. (2009). Population density in hydroponics culture for potato tuber-seed production (*Solanum tuberosum* L.). *Revista Chapingo, Serie Horticultura*, 15(3), 251-258.
- [17] Gericke, W. F. (1937). Hydroponics-crop production in liquid culture media. *Science*, 85(2198), 177-178.
- [18] Garlet, T. M. B., Santos, O. S., Medeiros, S. L. P., Garcia, D. C., Manfron, P. A., and Apel, M. A. (2007). Production of leaves, essential oil content, and quality of japanese mint (*Mentha arvensis* L. forma *piperascens holmes*) grown under hydroponics. *Revista Brasileira De Plantas Mediciniais*, 9(4), 72-79.
- [19] Girdthai, T., Jogloy, S., Kesmala, T., Vorasoot, N., Akkasaeng, C., Wongkaew, S., and Patanothai, A. (2010). Relationship between root characteristics of peanut in hydroponics and pot studies. *Crop Science*, 50(1), 159-167.
- [20] Guerrero-Lagunes, L. A., Del Mar Ruiz-Posadas, L., De Las Nieves Rodríguez-Mendoza, M., Soto-Hernández, M., and Castillo-Morales, A. (2011). Thyme hydroponics crops effect (*Thymus vulgaris* L.) on essential oil quality and yield. *Revista Chapingo, Serie Horticultura*, 17(2), 141-149.
- [21] Hoagland, D. R., and Arnon D. I. (1950). The water-culture method for growing plants without soil. *California Agricultural Experiment Station Circular*, 347,1-32.
- [22] Higashide, T., Kasahara, Y., Ibuki, T., and Sumikawa, O. (2005). Development of closed, energy saving hydroponics for sloping land. *Acta Horticulturae*, 691, 243-248
- [23] Hoque, M. M., Ajwa, H. A., and Smith, R. (2008). Nitrite and ammonium toxicity on lettuce grown under hydroponics. *Communications in Soil Science and Plant Analysis*, 39(1-2), 207-216.
- [24] Jensen, M. H. (1999). Hydroponics worldwide. *Acta Horticulturae*, 481, 719-729.
- [25] Kratky, B.A., and Bowen, J.E. (1988). Observation on a noncirculating hydroponic system for Tomato Production. *HortScience*, 23(5), 906-097.
- [26] Kanokwan, J., and Nobuo, N. (2008). Pharmacological aspects of *Andrographis paniculata* on health and its major diterpenoid constituent andrographolide. *Journal of Health Science*, 54(4), 370-381.
- [27] Koohakan, P., Jeanaksorn, T., and Nuntagij, I. (2008). Major diseases of lettuce grown by commercial nutrient film technique in Thailand. *KMITL Sci. Tech. J*, 8, 56-63.

- [28] Kitaya, Y., Hirai, H., Wei, X., Islam, A. F. M. S., and Yamamoto, M. (2008) Growth of sweet potato cultured in the newly designed hydroponic system for space farming. *Advances in Space Research*, 41, 730-735.
- [29] Kim, G., Hyun, D. Y., Kim, Y. O., Lee, S. E., Kwon, H., Cha, S. W., and Kim, Y. B. (2010). Investigation of ginsenosides in different parts of *Panax ginseng* cultured by hydroponics. *Korean Journal of Horticultural Science and Technology*, 28(2), 216-226.
- [30] Liew, O. W., Boey, W. S. L., Asundi, A. K., Chen, J. W., and He, D. M. (1999). Fibre optic spectrophotometry monitoring of plant nutrient deficiency under hydroponics culture conditions. *Proceedings of SPIE - the International Society for Optical Engineering*, 3740, 186-190.
- [31] Lopes, S. J., Dos Santos, P. M., Storck, L., Cocco, C., Damo, H. P., and Martini, L. F. D. (2007). Sampling chronology of hydroponics lettuce plants for adjusting vegetative growth curves. *Pesquisa Agropecuaria Brasileira*, 42(10), 1451-1457.
- [32] Lucena, J. J., Sentís, J. A., Villén, M., Lao, T., and Pérez-Sáez, M. (2008). IDHA chelates as a micronutrient source for green bean and tomato in fertigation and hydroponics. *Agronomy Journal*, 100(3), 813-818.
- [33] Laganowsky, A., Gomes, S. M., Whitelegge, J. P. and Nishio, J. N. (2009). Hydroponics on a chip: Analysis of the Fe deficient *Arabidopsis* thylakoid membrane proteome. *Journal of proteomics*, 72, 397-415.
- [34] Marin, A., and Dos Santos, D. M. M. (2008). Interaction of water stress and aluminum toxicity in pigeon pea cultivated in hydroponics. *Pesquisa Agropecuaria Brasileira*, 43(10), 1267-1275.
- [35] Mia, M. B., Shamsuddin, Z. H., Wahab, Z., and Marziah, M. (2010). Effect of plant growth promoting rhizobacterial (PGPR) inoculation on growth and nitrogen incorporation of tissue-cultured musa plantlets under nitrogen-free hydroponics condition. *Australian Journal of Crop Science*, 4(2), 85-90.
- [36] Martineau, V., Lefsrud, M., Naznin, M. T., and Kopsell, D. A. (2012). Comparison of light-emitting diode and high-pressure sodium light treatments for hydroponics growth of boston lettuce. *HortScience*, 47(4), 477-482.
- [37] Mayrapetyan, Kh. S., Alexanyan, J. S., Tadevosyan, A. H., and Mayrapetyan, S. Kh. (2012) Application of polymer as a means of water and nutrient elements saving in open-air hydroponics. *Acta Horticulturae*, 927, 935-940
- [38] Norton, G. J., Lou-Hing, D. E., Mehrag, A. A. and Price, A. H. (2008). Rice-arsenate interaction in hydroponics: whole genome transcriptional analysis. *Journal of Experimental Botany*, 59(8), 2267-2276.
- [39] Ortiz, A., Rotatori, H., Schreiber, E., and Von Roth, G. (2009). Hydroponic Farming in Mahasarakham: Integrating Hydroponics into the Agricultural Curriculum (Report). Mahasarakham University, 1-13
- [40] Ojha, S. K., Bharti, S., Joshi, S., Kumari, S., and Arya, D. S. (2012). Protective effect of hydroalcoholic extract of *Andrographis paniculata* on ischaemia-reperfusion induced myocardial injury in rats. *Indian J. Med. Res.*, 135, 414-421.

- [41] Paulus, D., Medeiros, S. L. P., Santos, O. S., Manfron, P. A., Paulus, E., and Fabbrin, E. (2007). Content and essential oil quality in mint (*Mentha arvensis* L.) produced in hydroponics and wield. *Revista Brasileira De Plantas Medicinai*s, 9(2), 80-87.
- [42] Park, J., and Kurata, K. (2009). Application of microbubbles to hydroponics solution promotes lettuce growth. *HortTechnology*, 19(1), 212-215.
- [43] Paz-Alberto and Sigua. (2013) Phytoremediation: A Green Technology to Remove Environmental Pollutants. *American Journal of Climate Change*, 2, 71-86.
- [44] Pezzarossa, B., Rosellini, I., Borghesi, E., Tonutti, P., and Malorgio, F. (2014). Effects of se-enrichment on yield, fruit composition and ripening of tomato (*solanum lycopersicum*) plants grown in hydroponics. *Scientia Horticulturae*, 165, 106-110.
- [45] Roberto, K. (2000). *How to Hydroponics Book*. 3<sup>rd</sup> Edition. ISN-0-9672026-0-4
- [46] Rababah, A., and Al-Shuha, A. (2009). Hydroponics reducing effluent's heavy metals discharge. *Water Science and Technology*, 59(1), 175-183.
- [47] Rodríguez-Delfín, A. (2012) Advances of hydroponics in Latin America. *Acta Horticulturae*, 947, Pages 23-32
- [48] Ruiz, M., and Taleisnik, E. (2013). Field hydroponics assessment of salt tolerance in *cenchrus ciliaris* (L.): Growth, yield, and maternal effect. *Crop and Pasture Science*, 64(6), 631-639.
- [49] Souret, F. F., and Weathers, P. J. (2000). The growth of saffron (*Crocus sativus* L.) in aeroponics and hydroponics. *Journal of Herbs, Spices and Medicinal Plants*, 7(3), 25-35.
- [50] Sigrimis, N., Arvanitis, K. G., Pasgianos, G. D., and Ferentinos, K. (2001). Hydroponics water management using adaptive scheduling with an on-line optimiser. *Computers and Electronics in Agriculture*, 31(1), 31-46.
- [51] Savvas, D. (2002). Automated replenishment of recycled greenhouse effluents with individual nutrients in hydroponics by means of two alternative models. *Biosystems Engineering*, 83(2), 225-236.
- [52] Smeets, K., Ruytinx, J., Belleghem, F.V., Semane, B., Lin, D., Vangronsveld, J. and Cuyper, A. (2008). Critical evaluation and statistical validation of a hydroponic culture system for *Arabidopsis thaliana*. *Plant Physiology and Biochemistry*, 46, 212-218.
- [53] Snow, A. M., and Ghaly, A. E. (2008). Use of barley for the purification of aquaculture wastewater in a hydroponics system. *American Journal of Environmental Sciences*, 4(2), 89-102.
- [54] Singh, G., Biswas, D. R., and Marwaha, T. S. (2010). Mobilization of potassium from waste mica by plant growth promoting rhizobacteria and its assimilation by maize (*Zea mays*) and wheat (*Triticum aestivum* L.): A hydroponics study under phytotron growth chamber. *Journal of Plant Nutrition*, 33(8), 1236-1251.
- [55] Samadi, A. (2011). Effect of particle size distribution of perlite and its mixture with organic substrates on cucumber in hydroponics system. *Journal of Agricultural Science and Technology*, 13(1), 121-129.

- [56] Sargsyan, E., Vardanyan, A., Ghalachyan, L., and Bulgadaryan, S. (2011). Cultivation of thymus by in vitro and hydroponics combined method. *World Academy of Science, Engineering and Technology*, 80, 129-132.
- [57] Shinohara, M., Aoyama, C., Fujiwara, K., Watanabe, A., Ohmori, H., Uehara, Y., and Takano, M. (2011). Microbial mineralization of organic nitrogen into nitrate to allow the use of organic fertilizer in hydroponics. *Soil Science and Plant Nutrition*, 57(2), 190-203.
- [58] Singh, D., Tiwari, A. and Gupta, R. (2012). Phytoremediation of lead from wastewater using aquatic plants. *Journal of Agricultural Technology*, 8(1), 1-11.
- [59] Sánchez-del-Castillo, F., González-Molina, L., Moreno-Pérez, E. C., Pineda-Pineda, J., and Reyes-González, C. E. (2014). Nutritional dynamics and yield of cucumber grown in hydroponics with and without recirculation of the nutrient solution. *Revista Fitotecnia Mexicana*, 37(3), 261-269.
- [60] Soares, T. M., Duarte, S. N., Silva, E. F. F., Melo, R. F., Jorge, C. A. and Oliveira, A. S. (2009). Experimental structure for evaluation of brackish water use in Lettuce hydroponic production. *Irriga. Botucatu*, 14(1), 102-114.
- [61] Thompson, J. L. C. (1978). Human Activities and Desert Expansion. *The Geographical Journal*, 144, 416-423.
- [62] Teragishi, A., Kanbara, Y., and Ono, H. (1998). Effects of low temperature storage, diameter of cuttings, and nutrient solution concentration on subsequent growth and fruit set of fig trees grown in hydroponics. *Journal of the Japanese Society for Horticultural Science*, 67(3), 386-390.
- [63] Teragishi, A., Kanbara, Y., and Ono, H. (2000). Effects of foliar application of choline chloride on the quality of winter-cropped, fig cv. masui-dauphine grown in hydroponics. *Journal of the Japanese Society for Horticultural Science*, 69(4), 390-395.
- [64] Tuberosa, R., Sanguineti, M. C., Landi, P., Giuliani, M. M., Salvi, S., and Conti, S. (2002). Identification of QTLs for root characteristics in maize grown in hydroponics and analysis of their overlap with QTLs for grain yield in the field at two water regimes. *Plant Molecular Biology*, 48(5), 697-712.
- [65] Tadevosyan, A. H., Mairapetyan, S. Kh., Buniatyan, R. J., Alexanyan, J. S., Galstyan, H. Kh., Stepanyan, B. T., and Davtyan, G. S. (2005) Humulus hydroponics in Armenia. *Acta Horticulturae*, 668, 235-240
- [66] Thomas, B. (2006). Light signals and flowering. *Journal of Experimental Botany*. 57(13), 3387-3393.
- [67] Ti-Da, G. E., Shi-Wei, S., Wu, J., Dong-Mei, T., and Dan-Feng, H. (2009). Influence of glycine-N concentration on the growth and nitrogen metabolism of tomato seedlings under sterile hydroponics cultivation. *Shengtai Xuebao Acta Ecologica Sinica*, 29(4), 1994-2002.
- [68] Trang, N. T. D., Schierup, H., and Brix, H. (2010). Leaf vegetables for use in integrated hydroponics and aquaculture systems: Effects of root flooding on growth, mineral composition and nutrient uptake. *African Journal of Biotechnology*, 9(27), 4186-4196.
- [69] Villagarcia, M. R., Carter Jr., T. E., Ruffy, T. W., Niewoehner, A. S., Jennette, M. W., and Arrellano, C. (2001). Genotypic rankings for aluminum tolerance of soybean roots grown in hydroponics and sand culture. *Crop Science*, 41(5), 1499-1507.

- [70] Von-Bieberstein, P., Xu, Y., Leslie Gunatilaka, A. A., and Gruener, R. (2014). Biomass production and withaferin a synthesis by *Withania somnifera* grown in aeroponics and hydroponics. *HortScience*, 49(12), 1506-1509.
- [71] Wang, X. F., and Ito, T. (1997). Effect of  $\text{NO}_3\text{-N}$  in the additional nutrient solution on the growth, yield and  $\text{NO}_3$  content in spinach plant grown in hydroponics. *Journal of the Japanese Society for Horticultural Science*, 66(2), 313-319.
- [72] Wang, Y. (1999). Growth and physiological characteristics of rice seedlings raised with long mat by hydroponics - comparison with young seedlings raised in soil. *Plant Production Science*, 2(2), 115-120.
- [73] Xing, Y. X., and Meng, X. D. (1999). Development and prospect of hydroponics in China. *Acta Horticultura*, 481, 753-760.
- [74] Yan, Q., Duan, Z., Mao, J., Li, X., and Dong, F. (2012) Effects of root-zone temperature and N, P, and K supplies on nutrient uptake of cucumber (*Curcumis sativus* L.) seedlings in hydroponics. *Soil Science and Plant Nutrition*, 58(6), 707-717
- [75] Zúñiga-Estrada, L., Martínez-Hernández, J. D. J., Baca-Castillo, G. A., Martínez-Garza, Á., Tirado-Torres, J. L., and Kohashi-Shibata, J. (2004). Bell pepper production in two irrigation systems under hydroponics conditions. *Agrociencia*, 38(2), 207-218 .
- [76] Zeng, Y., Guo, L., Huang, L., and Sun, Y. (2007). Plant hydroponics and its application prospect in medicinal plants study. *Zhongguo Zhongyao Zazhi*, 32(5), 374-376.